

Bakteriers känslighet för olika antibiotika (Del 1/2)

	Grampositiva aerob										Grampositiva anaerob	Gramnegativa aerob										G+ anaerob				
	S. aureus	Attihermylrykta streptokocker	Brakhamyloklar streptokocker (A, C, G)	Pneumokocker	E. faecalis	E. faecium	S. saprophyticus	Listeria	C. perfringens	C. difficile	Meningokocker	H. influenzae	Moraxella	E. coli	Klebsiella	Enterobacter	Proteus	Shigella, Salmonella	Campylobacter	Pseudomonas	Legionella	F. tularensis	Pasteurella	B. fragilis	Mycoplasma, Chlamydia	Borrelia
Fenoximetylpenicillin, penicillin V (oralt)												1, 2														
Bensylpenicillin, penicillin G (IV)												1, 2														
Pivmecillinam (oralt)							3							3	3	3, 4	3									
Amoxicillin (oralt) Ampicillin (IV)												2														
Amoxicillin/klavulansyra (oralt)																										
Flukloxacillin (oralt) Kloxacillin (IV)																										
Piperacillin/tazobaktam (IV)																4										
Cefadroxil (oralt)																4										
Ceftibuten (oralt)																4										
Cefuroxim (IV)																4										
Cefotaxim (IV)																4										
Ceftriaxon (IV)																4										1
Ceftazidim (IV)																4										1
Imipenem (IV)																										
Meropenem (IV)																										
Ertapenem (IV)																										
Aztreonam (IV)																4										

1: hög dos; 2: cirka 15-20 % resistenta; 3: endast nedre urinvägsinfektion; 4: inducerbar resistens; grå linje: osäker aktivitet.